

NEW ZEALAND MUDSNAIL

Potamopyrgus antipodarum

Photo courtesy of Larry Mayer (penny) and Daniel L. Gustafson (mudsnail), Minnesota Sea Grant..



The New Zealand mudsnail is a tiny freshwater snail that has established high densities in streams, rivers, and waterways all around the world. Since all introduced mudsnails in the United States reproduce asexually, it takes only one snail to start a new population.

SPECIES DESCRIPTION

New Zealand mudsnails are small, with adults ranging from 3-12 mm (0.1-0.5 in) in their native range, and 3.5-6 mm (0.1-0.2 in) in invasive populations. The shell is long, narrow, and coiled to the right in 7-8 whorls separated by deep grooves. Some morphs, such as those in the Great Lakes, can have either smooth shells, or shells with a keel in the middle of each whorl. The operculum, which covers the opening of the shell, is thin and ear-shaped. Often confused with other native snails, the New Zealand mudsnail is usually narrower, longer, and has more whorls than most native snails in the same genus.

NATIVE & INTRODUCED RANGES

Native to New Zealand, this species has spread to Europe, Asia, Australia, and North America. Two distinct populations exist in North America: a western population, which was discovered in the Snake River in Idaho and Wyoming in 1987; and a Great Lakes population, which was discovered in Lake Ontario and the St. Lawrence River in 1991. In 2007, New Zealand mudsnails were discovered in Lake Erie, about 6 km north of Presque Isle Bay in Erie County, Pennsylvania. Additional sampling in 2008 found New Zealand mudsnails in one small, unnamed stream near the Niagara River in New York State, and in 2011 they were found in an additional Lake Ontario tributary approximately 39 km from the first site. The most recent finding was in 2013 in Spring Creek in Center County, Pennsylvania; showing that they are expanding their range.

BIOLOGY & SPREAD

While it is thought that Western introductions occurred by stocking fish from overseas, the Great Lakes population most likely spread by contaminated ballast water from international ships. Once established, New Zealand mudsnails can reproduce very rapidly, producing an average of 230 juveniles per year. Although in its native New Zealand, both sexual and asexual populations exist, in its invasive range all populations are asexual, consisting of almost exclusively females, which can produce twice as many daughters as sexual females. New Zealand mudsnails can spread to new locations on recreational boating and fishing equipment, through the commercial movement of aquaculture products, and aquatic ornamental plants, in mud attached to the bills or legs of birds, or even within the gut of birds or fishes.






Photo courtesy of United States Geological Survey.



Map courtesy of United States Geological Survey.

NEW ZEALAND
 MUDSNAIL

NEW ZEALAND MUDSNAIL



Drawing courtesy of United States Geological Survey.

Sea Grant
Pennsylvania
www.paseagrant.org

Extension • Education • Research

Pennsylvania Sea Grant is a partnership of
The Pennsylvania State University,
The Commonwealth of Pennsylvania,
and NOAA.

Penn State is an affirmative action, equal
opportunity university.

 **pennsylvania**
DEPARTMENT OF ENVIRONMENTAL PROTECTION
COASTAL RESOURCES MANAGEMENT PROGRAM

 **Great Lakes RESTORATION**

Funded in part by PA DEP Coastal Resources
Management Program, The U.S. Fish and Wildlife
Service, and the Great Lakes Restoration Initiative

HABITAT

The New Zealand mudsnail flourishes in a variety of aquatic habitats, including springs, rivers, lakes, and estuaries. It has been found across a wide range of water temperatures (32-86°F [0-30°C]), substrates, water depths, productivity levels, and salinities. The diet of this snail is also variable, as it feeds on green algae and diatoms as well as detritus.

IMPACTS

Threat to Biodiversity

High population densities may allow New Zealand mudsnails to outcompete native fauna for food and substrate. In the Snake River, five species of native mollusks have been listed as endangered due in part to the establishment of the New Zealand mudsnail. Other studies have shown that in some rivers, New Zealand mudsnails make up over half the total invertebrate production, making them the dominate forage base for fish and other aquatic organisms. While they do not directly harm fish, their small hard shells may not be digestible, resulting in little to no nutritional value gained from eating the snails. Some species, such as trout, may avoid them as a food source altogether. New Zealand mudsnails can reach densities of up to 300,000 individuals/m² in some western U.S. rivers, dominating habitat and impacting preferred nesting sites.

PREVENTION & CONTROL

Know how to identify the New Zealand mudsnail. Always check for and remove plants, mud, and debris from boats, trailers, clothing, and equipment before leaving a water body. Drain all water from bait buckets, bilges, and live wells before transporting to new areas. Clean all gear and equipment with one of the following methods:

- 1.) Soak in hot water (60°C [140°F]) for at least five minutes (this method is not advised for Gore-tex®).
- 2.) Soak in a one-to-one solution of Formula 409® Cleaner Degreaser Disinfectant and water for at least five minutes then rinse with plain water. Simply spraying gear with the mixture does not work.
- 3.) Freeze for at least 6 hours (freezer temperature should be at -3°C [26°F] or below).

References:

- Alonso, A. and Castro-Díez, P. 2008. What Explains the invading success of the aquatic mud snail *Potamopyrgus antipodarum* (Hydrobiidae, Mollusca)? *Hydrobiologia*. 614: 107-116.
- California Department of Fish and Game. 2005. Controlling the Spread of New Zealand mud snails on wading gear. Office of Spill Prevention and Response Administrative Report 2005-02.
- Crosier, D. and Malloy, D. 2005. New Zealand Mudsnail (*Potamopyrgus antipodarum*). ANS Task Force website. < <http://anstaskforce.gov/spoc/nzms.php>>
- Levri, E.P, Kelly, A.A. and Love, E. 2007. The Invasive New Zealand Mud Snail (*Potamopyrgus antipodarum*) In Lake Erie. *Journal of Great Lakes Research*. 33(1): 1-6.
- Vinson, M. R. and Baker, M. A. 2008. Poor Growth of Rainbow Trout Fed New Zealand Mud Snails *Potamopyrgus antipodarum*. *North American Journal of Fisheries Management*. 28: 701-709.
- Levri, E.P., Colledge, E.D., Bilka, R. H., and Smith, B.J. 2012. The distribution of the invasive New Zealand mud snail (*Potamopyrgus antipodarum*) in streams in the lake Ontario and Lake Erie watersheds. *BioInvasions Records*. 1(3): 215-219.



Photo courtesy of Tim Worsfold,
World Register of Marine Species, EOL.

Photo courtesy of United States
Geological Survey.

