



Pennsylvania's Guide to  
**Identifying  
Baitfish**



# Acknowledgements

## Content Development

Kelly Donaldson

Amanda Mullen

Sara Stahlman

Amber Stilwell


Sarah Whitney

## Graphic Design

Jon Pelky, Graphics & Design, LLC

Pennsylvania Sea Grant is one of 34 Sea Grant programs supported by the National Sea Grant College Program and the National Oceanic and Atmospheric Administration.

Pennsylvania Sea Grant's mission is to enhance the use and conservation of Pennsylvania's coastal, marine, and Great Lakes resources to create a healthy environment, resilient and inclusive communities, and a strong and sustainable economy. Learn more online at <https://seagrant.psu.edu>.



---

Funding and support for this project was provided by the Great Lakes Restoration Initiative, U.S Environmental Protection Agency, National Oceanic and Atmospheric Administration, United States Fish and Wildlife Service, and the Pennsylvania Fish and Boat Commission.



# Table of Contents

Using This Guide .....	2
Aquatic Invasive Species Overview .....	3
Prevention Tips - Clean, Drain, Dry .....	6
Permitted Activities and Restrictions .....	8
Banned Species .....	9
<b>BAIT FISH ID</b>	
Fish Anatomy .....	10
<b>Family: Carps &amp; Minnows</b>	
Bighead Carp .....	12
Black Carp .....	13
Common Carp .....	14
Creek Chub .....	15
Emerald Shiner .....	16
European Rudd .....	17
Fallfish .....	18
Fathead Minnow .....	19
Golden Shiner .....	20
Grass Carp .....	21
Mimic Shiner .....	22
Northern Snakehead .....	23
Redside Dace .....	24
Rosyside Dace .....	25
Silver Carp .....	26
<b>Family: Catfish</b>	
Black Bullhead .....	27
<b>Family: Goby</b>	
Round Goby .....	28
Tubenose Goby .....	29
<b>Family: Herring</b>	
Alewife .....	30
Gizzard Shad .....	31
<b>Family: Livebearer</b>	
Eastern Mosquitofish .....	32
Western Mosquitofish .....	33
<b>Family: Topminnow</b>	
Eastern Banded Killifish .....	34
Mummichog .....	35
<b>Family: Perch</b>	
Banded Darter .....	36
Eurasian Ruffe .....	37
Greenside Darter .....	38
Iowa Darter .....	39
White Perch .....	40
<b>Family: Sucker</b>	
White Sucker .....	41
<b>Family: Sculpin</b>	
Sculpin .....	42
<b>Family: Stickleback</b>	
Brook Stickleback .....	43
Fourspine Stickleback .....	44
<b>Family: Sunfish</b>	
Bluegill .....	45
Glossary .....	46
Common Name Index .....	49
Scientific Name Index .....	50
Sources .....	51
Image Credits .....	56

# Using This Guide

This guide was developed to help bait shop owners, dealers, growers, harvesters, anglers, and others quickly identify invasive baitfish species.

These species can be mistakenly included in baitfish shipments, whether the bait is grown or collected. Knowing how to quickly identify and dispose of aquatic invasive species are vital skills in the effort to protect the Commonwealth's waters from their spread.

Invasive and native species can have very similar physical characteristics, making it difficult to recognize one versus the other. This guide provides quick ID details to help anyone identify invasive baitfish. Native baitfish are also featured in the guide.

Many invasive baitfish species originate from Asia or Europe. However, a species may be native to North America but not to Pennsylvania. Similarly, a species may be native to western Pennsylvania, but considered invasive in eastern Pennsylvania, and vice versa, so it's important to know which baitfish species are native, locally.

This guide also serves to help increase awareness about aquatic invasive species and the threats they pose to native species and Pennsylvania's waters.

## Invasive Fishbait

Fishbait such as crayfish, crabs, and the nymphs, larvae, and pupae of all insects (those spending any part of their life cycle in the water), can also be considered invasive.

## Anglers Support PA's Economy

**\$627.2 million** in economic output  
**7,110** jobs supported  
*(American Sportfishing Association)*

# Aquatic Invasive Species Overview

Aquatic invasive species (AIS) are aquatic animals and plants that have been introduced into water bodies where they do not live naturally. Here, they can cause detrimental ecological and economic impacts and/or harm to human health.

Global and domestic trade, recreational activities, and a changing climate are helping AIS spread at accelerated rates. As a result, AIS are changing the health and natural diversity of watersheds across the Commonwealth.

Preventing the introduction and spread of AIS, including baitfish, is critically important to protecting Pennsylvania's natural resources, and averting long-term ecosystem damage. Once AIS become widely established, controlling their spread is technically difficult, expensive, and eradication can be nearly impossible.

## Pennsylvania's Waters

**84K** miles of waterways

**77** miles of Lake Erie shoreline

**112** miles of Delaware Estuary coastline

**5** major watersheds

**\$40 Million**

**Economic Input to Lake Erie Community  
from Fishing Related Activities**

*(Dr. Alan Graefe, 2016)*

# Why It's Important to Stop the Spread of AIS

Baitfish dealers and anglers play a significant role in protecting the Commonwealth's waters and native species from the potential harm of AIS. Everyone's participation is needed to help stop their spread.

Aquatic invasive species can have detrimental impacts to ecosystems and native aquatic species. The *Baitfish Primer*, developed by Becky Cudmore and Nicholas Mandrak, describes several ways that AIS can impact native species, including:

- 1)** Aquatic invasive species can out-compete native species for food and may prey on them and their eggs. This reduces the abundance of native species and alters the natural food web of that ecosystem.
- 2)** Aquatic invasive species can cause habitat changes through behaviors like bottom-feeding and spawning which can destroy aquatic vegetation, increase turbidity, and negatively impact native species.
- 3)** Diseases and parasites can be transferred from aquatic invasive species to native species. Exposure to these diseases or parasites may lead to decreased abundance of native species.
- 4)** Native species are well adapted to their environment. Aquatic invasive species, not adapted to their new environment, may spawn with native species. Their offspring may look the same, but they are less adapted to their environment. Aquatic invasive individuals may also spawn with native individuals of closely related species. Their offspring (hybrids) may be less adapted to their environment or may be unable to reproduce. In most cases, spawning between introduced and native species will lead to a decreased abundance of native species. These impacts are not limited to aquatic invasive baitfishes.

# How are AIS introduced?

There are many ways AIS can be unintentionally spread.

## BOATING AND FISHING ACTIVITIES

**As Hitchhikers:** AIS can spread by attaching themselves to boats, trailers, equipment, fishing tackle, motors, clothing, diving gear, and boots.

**As Stowaways:** Microscopic AIS such as the Spiny Waterflea, Fishhook Waterflea, and Zebra Mussel larvae can be present in water in bait buckets, live wells, or bilges.

**As Discarded Bait:** AIS can be spread by discarding unused bait into a non-native waterbody. Additionally, baitfish shipments can accidentally contain non-native or invasive fish that can then be released into a new ecosystem.

## INTERNATIONAL SHIPPING

Species are transported in the ballast tanks of cargo ships and often released into non-native waters.


## AQUARIUM OWNERS

Without realizing the consequences, well-meaning individuals may release their aquarium pet into a waterbody when they are no longer able to care for it.

### CHECK YOUR GEAR!

Some AIS can survive out of the water on equipment or damp clothing for several days to weeks. Other AIS are so small they are very hard to see and therefore easily missed when inspecting equipment and gear.

New Zealand Mudsnail on a boot



# Prevention Tips - Clean, Drain, Dry

Preventing the introduction of aquatic invasive species to Pennsylvania's waters is the first and most vital step in stopping their spread. Anglers and recreational water users play a significant role in prevention efforts.

## CLEAN

Clean off visible aquatic plants, animals, and mud from watercraft, motor, trailer, waders, footwear, ropes, anchors, bait traps, dip nets, fishing lines, field gear, and any other equipment before leaving water access.

Scrub off any visible material on hull and footwear using a stiff brush. Rinse watercraft, trailer, and equipment with high pressure hot water when possible. Flush motor according to owner's manual.

**It is unlawful in Pennsylvania to transport plants, mud, or debris on boats and trailers.**

Boaters must also drain all water from their boat and remove drain/bilge plugs before leaving (Pennsylvania Regulations (58 Pa. Code Chapter 71a).

## CHECK THESE AREAS:




Diagram courtesy of the Aquatic Nuisance Species Taskforce's Recreational User Guidelines.

## DRAIN

Drain water from all equipment including watercraft, motor, bilge, bladder tanks, livewell, and portable bait containers before leaving water access.

## DRY

Dry everything for five days or more, unless otherwise required by local or state laws, when moving between waterbodies. Drying can kill small species not easily seen. If drying for 5 days isn't possible, wipe with a towel before reuse.

## DISPOSE

Drain water from all equipment  
Dispose of unwanted bait, fish parts, and packing materials in the trash; do not dump them in the water or on land.


Dumping any live bait into the water after fishing is illegal in Pennsylvania (58 Pa. Code Chapter 71a).

## Other Key Actions

- Use non-felt soled boots to further reduce the risk of spreading aquatic invasive species.
- Fish caught for eating or taxidermy should be cleaned at designated fish cleaning stations or placed on ice.

**Never move live fish or other organisms from one water body into another.**

## CHECK THESE AREAS:



# Permitted Activities and Restrictions

The following is a summary of guidance issued by the Pennsylvania Fish and Boat Commission related to baitfish. For a full listing of permitted activities and restrictions, see the Pennsylvania Fish and Boat Commission Fishing Summary Handbook published each year.

A person shall not

- Release or dispose of live baitfish into any water of this Commonwealth, except live baitfish baited on a hook for angling purposes.
- Use or possess Goldfish, Comets, Koi, and Common Carp as baitfish while fishing.
- Transfer a collected baitfish species from waters in this Commonwealth into another drainage of this Commonwealth where the species does not already exist.

A person may

- A person may collect live baitfish for angling.
- Use legally taken gamefish as bait.
- Use crayfish as bait if used in the water from which it was taken. Otherwise, the head must be immediately removed behind the eyes upon capture.


## Collecting, Purchasing, and Keeping Bait

- There is no closed season for collecting baitfish.
- There is no minimum size.
- The daily and possession limits are 50 combined species. Purchased fishbait or baitfish do not have a possession limit and do not count toward your daily limit of fishbait and baitfish taken from Commonwealth waterways. If fishing in a group, and possession limit exceeds 50, separate containers must be provided for the fishbait and baitfish.
- Retain sales receipt for purchased baitfish or fishbait. It serves as authorization for the possession for a period of 15 days.
- Stocked trout waters are closed for taking fishbait and baitfish from February 19 to the opening day of trout season. Most specially regulated trout waters are closed to taking fishbait and baitfish at all times.
- Collection of bait is permitted using rod and reel or handline, dip net, minnow seine, or minnow trap.
- Unattended minnow traps or baitfish containers must include the owner/user's name, address, and telephone number.

# Banned Species

It is illegal to possess, transport, sell, or introduce the following baitfish species, living or dead, in Pennsylvania.

- Bighead Carp (page 12)
- Black Carp (page 13)
- European Rudd (page 17)
- Round Goby (page 28)
- Eurasian Ruffe (page 37)
- Silver Carp (page 26)
- Snakehead (all species) (page 23)
- Tubenose Goby (page 29)



## DISPOSAL AND REPORTING

If you've found a suspected aquatic invasive species,


1. Use the "PA AIS" app to confirm.
2. Dispose of the species by freezing it.
3. Report it! Options to report include,
  - a. Call 1-833-Invasiv
  - b. Use the "PA AIS" app
  - c. Submit an online form to the Pennsylvania Fish & Boat Commission

**Scan Now!** To access AIS reporting methods and resources.


Or visit us online at <https://seagrant.psu.edu/aquatic-invasive-species/>




# Fish Anatomy



### Types of Caudal Fin



### Shapes of Homocercal Caudal Fins



## Bighead Carp

(*Hypophthalmichthys nobilis*)



**BANNED FOR SALE, BARTER, POSSESSION, OR  
TRANSPORTATION IN PENNSYLVANIA**

The Bighead Carp is native to eastern Asia. **It is considered invasive in Pennsylvania.** There is concern it could “stow away” in bait and be moved unintentionally. The Bighead Carp looks similar in appearance to the Silver Carp and Common Shiner.

---

### Quick ID:

**Distinctive Markings:** irregular gray-black blotches on sides

**Body:** broad, fusiform, laterally compressed

**Adult Length:** 39-58 in

**General Color:** dark gray top blending to white belly

**Eyes:** low set eyes, positioned downward

---

### Other Features:

**Head:** large

**Mouth:** large, terminal, protruding lower jaw

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** gray to dark gray in color, dorsal fin lacks spines

**Scales:** none on head, very small on body

## Black Carp

(*Mylopharyngodon piceus*)



**BANNED FOR SALE, BARTER, POSSESSION, OR  
TRANSPORTATION IN PENNSYLVANIA**

The Black Carp is native to eastern Asia from southern Russia to northern China. **It is considered invasive in Pennsylvania.** There are currently no reports of Black Carp in Pennsylvania. The Black Carp looks similar in appearance to the Grass Carp and the Common Carp.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** elongate, fusiform, slightly compressed

**Adult Length:** 39-58 in

**General Color:** dark brown to black on the back and sides, with some white on the underside

---

### Other Features:

**Head:** pointed

**Mouth:** small, terminal, lacks barbels

**Teeth or Barbels:** none

**Caudal Fin Shape:** large, forked

**Other Fin Notes:** fins are dark and lack spines

**Scales:** very large

## Common Carp

(*Cyprinus carpio*)



The Common Carp is native to Europe and Asia. **It is considered invasive in Pennsylvania.** It is illegal to use Common Carp as bait in Pennsylvania. The Common Carp looks similar in appearance to the Grass Carp.

---

### Quick ID:

**Distinctive Markings:** large scales outlined in black, two pairs of barbells on each side of the upper jaw

**Body:** heavy, stout

**Adult Length:** 16-31 in

**General Color:** olive brown to reddish brown, with yellow or yellow-white belly

---

### Other Features:

**Head:** short, rounded snout

**Mouth:** suckerlike, with two barbells on each side

**Teeth or Barbels:** barbells

**Caudal Fin Shape:** large, forked

**Other Fin Notes:** long dorsal fin, 17-23 soft rays

**Scales:** large, with dark edges and black spot at the base

## Creek Chub

(*Semotilus atromaculatus*)



The Creek Chub is native to most of the eastern United States. It is considered native in parts of Pennsylvania and is commonly used as live bait. The Creek Chub looks similar in appearance to the Fallfish.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** stout, robust, slightly compressed

**Adult Length:** 11 in

**General Color:** black or bluish with silvery belly

---

### Other Features:

**Head:** broad, blunt

**Mouth:** large, moderately oblique

**Teeth or Barbels:** small, single barbel in the corner of each jaw

**Caudal Fin Shape:** forked

**Other Fin Notes:** variable, brown to gray color, prominent dark spot on the base of the dorsal fin

**Scales:** small, appear cross-hatched

## Emerald Shiner

(*Notropis atherinoides*)



The Emerald Shiner is widely distributed across the United States, specifically in the Mississippi River Basin. **It is considered invasive in parts of Pennsylvania** and is commonly used as bait. The Emerald Shiner looks similar in appearance to other shiners.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** elongated, slender

**Adult Length:** 2-3 in

**General Color:** light olive to bluish green, sides are silvery, with a horizontal iridescent green or bluish green band the length of the body

**Eyes:** large

---

### Other Features:

**Head:** blunt snout

**Mouth:** moderately large, slightly upturned

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** dorsal fin behind pelvic fins, fins are clear

## European Rudd

(*Scardinius erythrophthalmus*)



### BANNED FOR SALE, BARTER, POSSESSION, OR TRANSPORTATION IN PENNSYLVANIA

The European Rudd is native to Europe and western Asia. **It is considered invasive in Pennsylvania.** There is concern it could be a bait stowaway. The European Rudd looks similar in appearance to the Common Shiner.

---

#### Quick ID:

**Distinctive Markings:** beginning of dorsal fin set well behind the front of the pelvic fin

**Body:** stocky, robust, elliptical

**Adult Length:** 19 in

**General Color:** brownish green on top, brassy yellow to roseay on sides, fading to silver underneath.

**Eyes:** iris is orange to yellow

---

#### Other Features:

**Head:** small

**Mouth:** terminal, lower lip is sharply angled with protruding lower jaw

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** pectoral, pelvic, and anal fins are a distinct red orange color

**Scales:** large

## Fallfish

(*Semotilus corporalis*)



The Fallfish is native to the northeast United States. **It is native in parts of Pennsylvania and invasive in others.** Its spread in Pennsylvania has been attributed to bait release. The Fallfish looks similar in appearance to the Creek Chub.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** moderately compressed

**Adult Length:** 20 in

**General Color:** olive to golden brown

---

### Other Features:

**Head:** rounded snout

**Mouth:** large, overhanging

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** fins are gray to brown

**Scales:** dorsal scales are darkly outlined

# FAMILY - CARPS & MINNOWS

## Fathead Minnow

(*Pimephales promelas*)



The Fathead Minnow is **native to much of the central United States and northwestern Pennsylvania; however, it is considered invasive in other parts of Pennsylvania**. It is commonly used as live bait.

- Breeding males acquire a large, gray fleshy growth on the nape, and approximately 16 white breeding tubercles on the snout.

---

### Quick ID:

**Distinctive Markings:** spot at base of tail fin

**Body:** chubby, moderately compressed

**Adult Length:** 3 in

**General Color:** dark olive or brown, sides are dusky, fading to pale belly

---

### Other Features:

**Head:** blunt and round snout

**Mouth:** slightly subterminal

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** short, rounded fins that are clear to gray or brown

**Scales:** scales ahead of dorsal fin are small and crowded

## Golden Shiner

(*Notemigonus crysoleucas*)



The Golden Shiner is native to the entire eastern half of the United States. **It is not considered invasive in Pennsylvania.** It is commonly used as live bait. The Golden Shiner looks similar in appearance to other shiners.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** extremely compressed

**Adult Length:** average 2-4 in, large 8-10 in

**General Color:** dark olive green, sides may be golden, brassy, or silvery

---

### Other Features:

**Head:** pointed snout

**Mouth:** small, upturned

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** anal fin is gray with long base

**Scales:** large

## Grass Carp

(*Ctenopharyngodon idella*)



The Grass Carp is native to eastern Asia including China and Russia. **It is considered invasive in Pennsylvania.** In Pennsylvania, a permit is required to stock triploid Grass Carp, a sterile form that is unable to reproduce. Stocking is only permitted in ponds to prevent the spread through waterways and river systems. The Grass Carp looks similar in appearance to the Common Carp.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** oblong, slender

**Adult Length:** 24-49 in

**General Color:** olive to silvery white, fins are clear to gray brown

---

### Other Features:

**Head:** slightly flattened

**Mouth:** small, terminal

**Teeth or Barbels:** none

**Caudal Fin Shape:** large, forked

**Other Fin Notes:** dorsal fin has three simple rays and seven branched rays

**Scales:** large, dark edges and black spot at the base

## Mimic Shiner

(*Notropis volucellus*)



The Mimic Shiner is native from Vermont south to Texas and the Gulf of Mexico, and north to the Great Lakes. **It is considered invasive in Pennsylvania.** Its introduction in Pennsylvania has been attributed to bait release. The Mimic Shiner looks similar in appearance to darters and gobies.

---

### Quick ID:

**Distinctive Markings:** dark pigmentation around anus, base of anal fin

**Body:** laterally compressed

**Adult Length:** 2-3 in

**General Color:** yellowish olive, silvery sides, white belly

**Eyes:** large

---

### Other Features:

**Head:** broad, round snout

**Mouth:** slightly upturned

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** clear fins

**Scales:** increase in size from front to back, upper sides are dark edged

## Northern Snakehead

(*Channa argus*)



**BANNED FOR SALE, BARTER, POSSESSION, OR  
TRANSPORTATION IN PENNSYLVANIA**

The Northern Snakehead is native to China, Russia, and Korea.

**It is considered invasive in Pennsylvania.** Its introduction in Pennsylvania has been attributed to pet release by aquarists and as bait by anglers. It looks similar in appearance to the native Bowfin and the native and endangered Burbot.

---

### Quick ID:

**Distinctive Markings:** snake-shaped head, truncated caudal fin, long anal fin

**Body:** cylindrical, long, thin

**Adult Length:** 33 in

**General Color:** tan with dark brown mottling

---

### Other Features:

**Head:** snake-shaped, flattened

**Mouth:** large

**Teeth or Barbels:** canine-like, sharp, many

**Caudal Fin Shape:** truncated

**Other Fin Notes:** long dorsal and anal fins

**Scales:** present on top of head

## Redside Dace

(*Clinostomus elongatus*)



The Redside Dace is native to the Upper Susquehanna River Basin, and the Great Lakes. **It is not considered invasive in Pennsylvania.** The Redside Dace looks similar in appearance to the Rosy side Dace.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** laterally compressed

**Adult Length:** 4 in

**General Color:** dark back with red band on lower side

---

### Other Features:

**Head:** pointed snout

**Mouth:** large, oblique

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** fins are light brown to clear

**Scales:** small

## Rosyside Dace

(*Clinostomus funduloides*)



The Rosyside Dace is native to the Atlantic Slope from the Lower Delaware River in Pennsylvania south to Georgia and west to southern Ohio, Tennessee, and Kentucky. **It is not considered invasive in Pennsylvania.** The Rosyside Dace looks similar in appearance to the Redside Dace.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** laterally compressed

**Adult Length:** 4 in

**General Color:** olive fading to white

---

### Other Features:

**Head:** pointed snout

**Mouth:** large, oblique

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** fins are olive to brown

**Scales:** small

## Silver Carp

(*Hypophthalmichthys molitrix*)



**BANNED FOR SALE, BARTER, POSSESSION, OR  
TRANSPORTATION IN PENNSYLVANIA**

The Silver Carp is native to eastern Asia. **It is considered invasive in Pennsylvania.** Resource managers are concerned that it may spread as a bait stowaway. The Silver Carp looks similar in appearance to the Bighead Carp and the Common Shiner.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** laterally compressed

**Adult Length:** 16-28 in

**General Color:** olive to grayish black, silvery sides blending to white belly

**Eyes:** large, low on head, more forward-facing

---

### Other Features:

**Head:** large

**Mouth:** large and terminal, lower jaw slightly protruding

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked


**Other Fin Notes:** short dorsal fin, lacks spines, fins are gray to dark

**Scales:** very small

# FAMILY - CATFISH

## Black Bullhead

(*Ameiurus melas*)



The Black Bullhead is native to the Great Lakes and Mississippi River basins. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** There is concern it could stow away in bait and be unintentionally moved. The Black Bullhead looks similar in appearance to the Brown Bullhead and the Yellow Bullhead.

---

### Quick ID:

**Distinctive Markings:** has adipose fin

**Body:** stout

**Adult Length:** 24 in

**General Color:** dark brown to gray

---

### Other Features:

**Head:** variable

**Mouth:** terminal mouth, upper lip extends past bottom

**Teeth or Barbels:** present, dark, whisker-like

**Adipose Fin:** Yes

**Caudal Fin Shape:** rounded, slightly notched

**Other Fin Notes:** fin coloration is similar to body

**Scales:** none

## Round Goby

(*Neogobius melanostomus*)



**BANNED FOR SALE, BARTER, POSSESSION, OR  
TRANSPORTATION IN PENNSYLVANIA**

The Round Goby is native to Eurasia. **It is considered invasive in Pennsylvania.** It was introduced in the United States through ballast water and has spread through bait release or by stowing away in bait buckets. The Round Goby looks similar in appearance to the Tubenose Goby, darters, and sculpins.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** cylindrical, somewhat flat underside

**Adult Length:** 3-10 in

**General Color:** solid slate gray, can be mottled with olive green, black, gray, and brown spots. Males can be nearly solid black.

**Eyes:** frog-like

---

### Other Features:

**Head:** large, round

**Mouth:** wide, large lips

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** black spot on the dorsal fin, light brown to clear fins

**Scales:** all over

# FAMILY - GOBY

## Tubenose Goby (*Proterorhinus semilunaris*)



### BANNED FOR SALE, BARTER, POSSESSION, OR TRANSPORTATION IN PENNSYLVANIA

The Tubenose Goby is native to the Black and Caspian seas in Europe. **It is considered invasive in Pennsylvania.** It was introduced in the United States through ballast water and can be spread through bait release or by stowing away in bait buckets. The Tubenose Goby looks similar in appearance to the Round Goby and sculpins.

---

#### Quick ID:

**Distinctive Markings:** has fused pelvic fins

**Body:** cylindrical body, somewhat flat underside

**Adult Length:** 2-4 in

**General Color:** light brown with dark brown blotches that may form vertical bars.

---

#### Other Features:

**Head:** blunt or rounded snout

**Mouth:** wide, large lips

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** triangular black spot is present at base of caudal fin followed by two white spots

**Scales:** small

# FAMILY - HERRING

## Alewife

(*Alosa pseudoharengus*)



The Alewife is native to the Atlantic Coast of the United States. **It is invasive in parts of Pennsylvania and native in other parts of Pennsylvania.** It is commonly used as live bait. The Alewife looks similar in appearance to the Gizzard Shad.

---

### Quick ID:

**Distinctive Markings:** black spot on each side just behind head

**Body:** small, slender, laterally compressed

**Adult Length:** 10-15 in

**General Color:** silver (bluish to green), often has coppery sheen, with horizontal darker stripes

**Eyes:** relatively large, obvious eyelid

---

### Other Features:

**Head:** broad and triangular

**Mouth:** superior

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** fins are brownish gray

**Scales:** row of scales form sharp edge along midline of the belly

# FAMILY - HERRING

## Gizzard Shad

(*Dorosoma cepedianum*)



The Gizzard Shad is native to the southern Great Lakes region, and from Pennsylvania to Florida and Texas. **It is considered native in parts of Pennsylvania and invasive in other parts of Pennsylvania.** It is commonly used as live bait. The Gizzard Shad looks similar in appearance to the Alewife.

---

### Quick ID:

**Distinctive Markings:** closely packed gill rakers

**Body:** short, moderately to strongly compressed

**Adult Length:** 20 in

**General Color:** dark on top fading to silvery white belly

**Eyes:** has eyelids

---

### Other Features:

**Head:** blunt snout

**Mouth:** small, subterminal to inferior

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** long trailing last dorsal ray, brown to gray fins

**Scales:** none on head, scales on body are thin

## Eastern Mosquitofish

(*Gambusia holbrooki*)



The Eastern Mosquitofish is native to the Atlantic and Gulf Slope drainages. **It is considered invasive in Pennsylvania.** It has been observed as a stowaway in bait shipments. The Eastern Mosquitofish looks similar in appearance to the Western Mosquitofish.

- Females usually have a black stripe near the eye area but are more easily recognized by a black spot on their belly called the gravid spot which darkens during pregnancy.

---

### Quick ID:

**Distinctive Markings:** dark teardrop shape below eyes

**Body:** compressed, females have rounded belly

**Adult Length:** 2 in

**General Color:** olive gray to yellowish

---

### Other Features:

**Head:** flattened

**Mouth:** small, upturned

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** dusky to black spots on dorsal and caudal fins, small dorsal fin starting behind anal fin, clear fins

**Scales:** darkly outlined

## Western Mosquitofish (*Gambusia affinis*)



The Western Mosquitofish is native to the Atlantic and Gulf Slope drainages from New Jersey to New Mexico and Illinois to Texas. **It is considered invasive in Pennsylvania.** Its introduction in Pennsylvania has been attributed to bait release. The Western Mosquitofish looks similar in appearance to the Eastern Mosquitofish.

- Females are generally larger than males. During pregnancy, females tend to have a distended abdomen with a black spot called a gravid spot which darkens during pregnancy.

---

### Quick ID:

**Distinctive Markings:** small teardrop streaks below eyes

**Body:** short

**Adult Length:** 2-3 in

**General Color:** dull gray or brown

---

### Other Features:

**Head:** flattened

**Mouth:** pointed upwards

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** fins are clear, small light speckling on caudal and dorsal fins

**Scales:** lightly outlined in black on back and sides

# FAMILY - TOPMINNOW

## Eastern Banded Killifish (*Fundulus diaphanus*)



The Eastern Banded Killifish is native in the Atlantic Slope drainages of the northeast United States to South Carolina and west to the Dakotas. **It is native in some parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its introduction in Pennsylvania has been attributed to bait release. The Eastern Banded Killifish looks similar in appearance to darters and gobies.

- Females have fewer than 9 thin, distinctly black bands that often do not span the full width of the body. Males have more than 9 silvery bands which are less distinct and closer together.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** Laterally compressed

**Adult Length:** 2-4 in

**General Color:** dark gray on the back fading around sides to white or yellow belly. Spawning males have yellow fins.

---

### Other Features:

**Head:** flattened, bluntly pointed snout

**Mouth:** small, slightly upturned

**Teeth or Barbels:** small teeth on both jaws

**Caudal Fin Shape:** rounded to truncated

**Other Fin Notes:** clear to gray fins

**Scales:** small, all over

# FAMILY - TOPMINNOW

## Mummichog

(*Fundulus heteroclitus*)



The Mummichog is native to marine, brackish, and occasionally freshwater areas along the Atlantic Coast from Maine to northeastern Florida. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its spread in Pennsylvania has been attributed to bait release. The Mummichog looks similar in appearance to shiners.

- Males typically present with bright yellow fins, and females are larger in body size.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** stout

**Adult Length:** 1-3 in

General Color: brownish green to green or olive

---

### Other Features:

**Head:** flat, short blunt snout

**Mouth:** lower lip juts out

**Teeth or Barbels:** teeth

**Caudal Fin Shape:** rounded or squared off

**Other Fin Notes:** females have green tinted dorsal and anal fins, males have yellow pectoral and anal fins

**Scales:** all over

## Banded Darter

(*Etheostoma zonale*)



The Banded Darter is native to Lake Michigan and the Mississippi River basins from New York to Minnesota, and south to northern Georgia, northern Alabama, and southern Arkansas. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its spread in Pennsylvania has been attributed to bait release. The Banded Darter looks similar in appearance to other darters and gobies.

- Males possess more striking colors during spawning while females maintain neutral earth tones.

---

### Quick ID:

**Distinctive Markings:** dark green vertical bands along the body, and 3 to 4 dark spots on the base of the tail fin

**Body:** short and stubby

**Adult Length:** 3 inches

**General Color:** green and brown, belly is yellow to white

---

### Other Features:

**Head:** rounded snout

**Mouth:** subterminal, upper lip is connected to snout by a narrow ridge of skin

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** all fins have narrow dark lines, the dorsal fins have a red to rust colored band at the base

**Scales:** all over

# FAMILY - PERCH

## Eurasian Ruffe

(*Gymnocephalus cernua*)



**BANNED FOR SALE, BARTER, POSSESSION, OR  
TRANSPORTATION IN PENNSYLVANIA**

The Eurasian Ruffe is native to Eurasia. **It is considered invasive in Pennsylvania.** Its introduction in Pennsylvania has been attributed to bait release. The Eurasian Ruffe looks similar in appearance to the Yellow Perch and the White Perch.

---

### Quick ID:

**Distinctive Markings:** rows of dark spots between spines on dorsal fin, very slimy when handled, sharp spines on gill covers

**Body:** slender, slightly compressed

**Adult Length:** 4-6 in

**General Color:** greenish to brown on top with dark patches on lighter brown sides, yellow belly has dark spots

**Eyes:** glassy

---

### Other Features:

**Head:** large

**Mouth:** small, downturned

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** dorsal fin behind pelvic fins, fins are clear to yellow

**Scales:** on body, none on head

# FAMILY - PERCH

## Greenside Darter (*Etheostoma blennioides*)



The Greenside Darter is native to the Great Lakes and Mississippi River basins, and from New York to Maryland to Kansas and Oklahoma. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its spread in Pennsylvania has been attributed to bait release. The Greenside Darter looks similar in appearance to other darters.

- During spawning season, males display striking colors while females maintain neutral earth tones.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** cylindrical

**Adult Length:** 2-3 in

**General Color:** brownish to green, dark or reddish spots present on the top of back, green bands along sides

**Eyes:** large

---

### Other Features:

**Head:** rounded snout

**Mouth Shape:** small, terminal

**Teeth or Barbels:** none

**Caudal Fin Shape:** slightly notched

**Other Fin Notes:** large well-developed pectoral fins, and green dorsal fins with reddish bands at the base

**Scales:** all over

# FAMILY - PERCH

## Iowa Darter

(*Etheostoma exile*)



### REGULATED AS ENDANGERED IN PENNSYLVANIA

The Iowa Darter is native to the northern central United States. **It is native to parts of Pennsylvania and is not considered invasive in Pennsylvania.** It has been observed as a stowaway in bait shipments. The Iowa Darter looks similar in appearance to other darters.

- During spawning, males display striking colors while females maintain neutral earth tones.

#### Quick ID:

**Distinctive Markings:** teardrop mark under eyes, 9 to 12 dark blotches on sides

**Body:** slender

**Adult Length:** 2-3 in

**General Color:** greenish or brownish

**Eyes:** large

#### Other Features:

**Head:** rounded snout

**Mouth:** terminal

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** front dorsal fin is green with red stripe

**Scales:** all over

## White Perch

*(Morone americana)*



The White Perch is native to the Atlantic Coast of the United States. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its spread in Pennsylvania has been attributed to bait release. The White Perch looks similar in appearance to the White Bass and the Yellow Perch.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** laterally compressed

**Adult Length:** 5-7 in

**General Color:** silvery gray to greenish brown fading to silvery white

---

### Other Features:

**Head:** steep slope from nose to dorsal fin

**Mouth:** large, subterminal, extends just below the eye

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** dorsal fin has spiny and soft portions separated by a deep notch, fins are gray to brown

**Scales:** all over

# FAMILY - SUCKER

## White Sucker

(*Catostomus commersonii*)



The White Sucker is native to the Atlantic Coast, Arctic, Great Lakes, and the Mississippi River basin. **It is not considered invasive in Pennsylvania.** The White Sucker looks similar in appearance to the Common Carp.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** round

**Adult Length:** 25 in

**General Color:** olive brown to black fading to white on belly

---

### Other Features:

**Head:** rounded snout

**Mouth:** inferior, faces downward

**Teeth or Barbels:** none

**Caudal Fin Shape:** forked

**Other Fin Notes:** none

**Scales:** increase in size from front to back

## Sculpin

(*Cottus spp.*)



The Sculpin is native to parts of the eastern and western United States. **It is not considered invasive in Pennsylvania.** The Sculpin looks similar in appearance to the Round Goby and the Tubenose Goby.

---

### Quick ID:

**Distinctive Markings:** none

**Body:** cylindrical, somewhat flat underside

**Adult Length:** 6 in

**General Color:** brown, may have dark blotches that appear as vertical bands

**Eyes:** large

---

### Other Features:

**Head:** large, rounded

**Mouth:** large and wide with large lips

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** large, pointy fins that are brown to clear

**Scales:** none

# FAMILY - STICKLEBACK

## Brook Stickleback

(*Culaea inconstans*)



The Brook Stickleback is native to the Great Lakes and Mississippi River basins from Vermont to Montana, and south to Ohio and Illinois. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** It has been observed as a stowaway in bait shipments and may have been introduced via bait bucket releases. The Brook Stickleback looks similar in appearance to the Fourspine Stickleback.

---

### Quick ID:

**Distinctive Markings:** 4 to 6 dorsal spines on back

**Body:** deep, compressed

**Adult Length:** 3 in

**General Color:** olive, belly tends to be silver white or light green, may have pale vertical lines

---

### Other Features:

**Head:** conical

**Mouth:** lower jaw extends past upper

**Teeth or Barbels:** sharp teeth on both jaws

**Caudal Fin Shape:** rounded

**Other Fin Notes:** fins are clear to olive

**Scales:** none, has small bony plates along sides

# FAMILY - STICKLEBACK

## Fourspine Stickleback

(*Apeltes quadratus*)



The Fourspine Stickleback is native to Europe and the North American Atlantic Coast from New Brunswick to North Carolina. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its spread in Pennsylvania has been attributed to bait release. The Fourspine Stickleback looks similar in appearance to the Brook Stickleback.

---

### Quick ID:

**Distinctive Markings:** 2 to 4 free dorsal spines close to each other, has a bony ridge on each side of abdomen

**Body:** triangular, flat belly, sharp back

**Adult Length:** 2 in

**General Color:** brown on back and sides with white belly, body covered in dark mottling

---

### Other Features:

**Head:** tapers to pointed nose

**Mouth:** terminal, slightly upturned

**Teeth or Barbels:** none

**Caudal Fin Shape:** rounded

**Other Fin Notes:** dorsal and anal fins are rounded, brownish white, slim base at tail, clear pelvic fins are sometimes reddish

**Scales:** on sides

## Bluegill

(*Lepomis macrochirus*)



The Bluegill is native to the St. Lawrence, Great Lakes, and Mississippi River Basins from New York to Minnesota and south to the Gulf of Mexico. **It is native in parts of Pennsylvania and considered invasive in other parts of Pennsylvania.** Its introduction in Pennsylvania has been attributed to bait release. Juvenile Bluegill are similar in appearance to darters and gobies.

---

### Quick ID:

**Distinctive Markings:** flexible black ear flap

**Body:** compressed

**Adult Length:** 9 in

**General Color:** blue green back blends to white belly, with 5 to 9 dark vertical bands on sides which lighten

**Eyes:** large, dark

---

### Other Features:

**Head:** bluntly pointed

**Mouth:** terminal, small

**Teeth or Barbels:** none

**Caudal Fin Shape:** shallow fork with rounded tips

**Other Fin Notes:** two spiny-ray dorsal fins are continuous, dusky rounded spot on rear dorsal fin

**Scales:** all over

# Glossary

**Anal Fin:** a median unpaired fin on the lower posterior part of the body behind the vent and sometimes confluent with the caudal fin

**Anterior:** situated before or toward the front

**Aquatic Invasive Species:** freshwater or marine organism that has spread or been introduced beyond its native range and is either causing harm or has the potential to cause harm

**Baitfish:** small marine or freshwater fish that are used as bait to attract larger predatory fish

**Barbel:** whisker like sensory organ near the mouth

**Bilge:** outer surface of a ship's hull where the bottom curves to meet the vertical sides

**Caudal:** directed towards the hind part of the body

**Caudal Fin:** the tail fin, located at the end of the caudal peduncle; used for propulsion

**Caudal Peduncle:** tapered region behind the dorsal and anal fins where the caudal fin attaches to the body

**Compressed:** having a narrow shape as if flattened

- **Laterally Compressed:** flattened from side to side

**Conical:** resembling a cone especially in shape

**Cylindrical:** relating to or having the form or properties of a cylinder

**Depressed:** flattened from top to bottom

**Diphyceral:** caudal tail shape; vertebrae extend to the tip of the tail and the tail is symmetrical and expanded

**Dorsal:** located on the top of a fish, used for stabilizing in the water and directing through turns and stops

**Dorsal Fin:** fish may have three, located on the back, serves as protection against rolling and assists in sudden turns and stops

**Elliptical:** of, relating to, or shaped like an ellipse

**Elongate:** to extend the length of

**Emarginate:** having the margin notched

**Fishbait:** all insects spending any part of their life cycle in the water

**Forked:** resembling a fork especially in having one end divided into two or more branches or points

**Fusiform:** tapering toward each end

**Gill Raker:** any of the bony processes on a gill arch that divert solid substances away from the gills

**Heterocercal:** of a fish tail fin: having the upper lobe larger than the lower with the vertebral column extending into the upper lobe

**Homocercal:** of a fish tail fin: having the upper and lower lobes approximately symmetrical and the vertebral column ending at or near the middle of the base

**Inferior:** situated in a relatively low posterior or ventral position in a quadrupedal body

**Lateral Line:** a canal along the side of a fish containing pores that open into tubes supplied with sense organs sensitive to low vibrations

**Livewell:** a well for keeping fish alive in a fishing boat by allowing water to circulate through it

**Lunate:** shaped like a crescent

**Oblique:** neither perpendicular nor parallel

# Glossary

**Operculum:** the covering of the gills of a fish

**Pectoral Fin:** either of the fins of a fish that correspond to the forelimbs of a quadruped

**Pelvic Fin:** one of the paired fins of a fish that are homologous with the hind limbs of a quadruped

**Peduncle:** a stalk-like part by which an organ is attached to an animal's body, or by which a barnacle or other sedentary animal is attached to a substrate

**Posterior:** situated behind

**Ray:** compound, segmented, and branched bony structures which support a fin

**Rounded:** made round: flowing rather than jagged or angular

**Superior:** situated higher up

**Terminal:** of or relating to an end, extremity, boundary, or terminus, a terminal pillar, growing at the end of a branch or stem

- **Subterminal:** situated or occurring near but not precisely at an end

**Truncate:** Appearing to end abruptly, shortened, cut off at the end

**Tubercles:** a small knobby prominence or excrescence especially on a plant or animal

**Ventral:** being or located near or on the anterior or lower surface of an animal opposite the back

## Common Name Index

Alewife .....	30
Banded Darter .....	36
Bighead Carp .....	12
Black Bullhead .....	27
Black Carp.....	13
Bluegill.....	45
Brook Stickleback .....	43
Common Carp.....	14
Creek Chub.....	15
Eastern Banded Killifish.....	34
Eastern Mosquitofish.....	32
Emerald Shiner .....	16
Eurasian Ruffe .....	37
European Rudd.....	17
Fallfish .....	18
Fathead Minnow.....	19
Fourspine Stickleback .....	44
Gizzard Shad .....	31
Golden Shiner .....	20
Grass Carp .....	21
Greenside Darter.....	38
Iowa Darter .....	39
Mimic Shiner .....	22
Mummichog .....	35
Northern Snakehead.....	23
Redside Dace .....	24
Rosyside Dace .....	25
Round Goby.....	28
Sculpin.....	42
Silver Carp.....	26
Tubenose Goby .....	29
Western Mosquitofish .....	33
White Perch .....	40
White Sucker .....	41

# Scientific Name Index

<i>Alosa pseudoharengus</i> .....	30
<i>Ameiurus melas</i> .....	27
<i>Apeltes quadracus</i> .....	44
<i>Catostomus commersonii</i> .....	41
<i>Channa argus</i> .....	23
<i>Clinostomus elongatus</i> .....	24
<i>Clinostomus funduloides</i> .....	25
<i>Cottus spp.</i> .....	42
<i>Ctenopharyngodon idella</i> .....	21
<i>Culæa inconstans</i> .....	43
<i>Cyprinus carpio</i> .....	14
<i>Dorosoma cepedianum</i> .....	31
<i>Etheostoma blennioides</i> .....	38
<i>Etheostoma exile</i> .....	39
<i>Etheostoma zonale</i> .....	36
<i>Fundulus diaphanus</i> .....	34
<i>Fundulus heteroclitus</i> .....	35
<i>Gambusia affinis</i> .....	33
<i>Gambusia holbrooki</i> .....	32
<i>Gymnocephalus cernua</i> .....	37
<i>Hypophthalmichthys molitrix</i> .....	26
<i>Hypophthalmichthys nobilis</i> .....	12
<i>Lepomis macrochirus</i> .....	45
<i>Morone americana</i> .....	40
<i>Mylopharyngodon piceus</i> .....	13
<i>Neogobius melanostomus</i> .....	28
<i>Notemigonus crysoleucas</i> .....	20
<i>Notropis atherinoides</i> .....	16
<i>Notropis volucellus</i> .....	22
<i>Pimephales promelas</i> .....	19
<i>Proterorhinus semilunaris</i> .....	29
<i>Scardinius erythrophthalmus</i> .....	17
<i>Semotilus atromaculatus</i> .....	15
<i>Semotilus corporalis</i> .....	18

## Sources

**Alewife** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=490>

**Banded Darter** – Illinois Department of Natural Resources, Wild About Illinois Fishes! <https://dnr.illinois.gov/education/wildaboutpages/wildaboutfishes/wafperch/wafbandeddarter.html>; Oklahoma Department of Wildlife Conservation, Oklahoma Field Guide: <https://www.wildlifedepartment.com/wildlife/field-guide/fish/banded-darter>; United States Geologic Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=818>

**Bighead Carp** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=551>

**Black Bullhead** – North Dakota Game and Fish Department, Common and Species of Conservation Priority Fish in North Dakota: <https://gf.nd.gov/wildlife/id/black-bullhead>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=730>

**Black Carp** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=573>

**Bluegill** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=385>; Wisconsin Department of Natural Resources: [https://dnr.wisconsin.gov/sites/default/files/topic/Fishing/Species\\_bluegill.pdf](https://dnr.wisconsin.gov/sites/default/files/topic/Fishing/Species_bluegill.pdf)

**Brook Stickleback** – The Regents of the University of Michigan: [https://www.biokids.umich.edu/critters/Culaea\\_inconstans/](https://www.biokids.umich.edu/critters/Culaea_inconstans/); United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=701>

## Sources

**Creek Chub** – Connecticut Department of Energy and Environmental Protection: <https://portal.ct.gov/DEEP/Fishing/Freshwater/Freshwater-Fishes-of-Connecticut/Creek-Chub>; Texas State University, San Marcos, Department of Biology. Freshwater Fishes: <http://txstate.fishesoftexas.org/semotilus%20atromaculatus.htm>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=649>

**Common Carp** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=4>

**Eastern Banded Killifish** – Connecticut Department of Energy and Environmental Protection: <https://portal.ct.gov/DEEP/Fishing/Freshwater/Freshwater-Fishes-of-Connecticut/Banded-Killifish>; Iowa Department of Natural Resources: <https://programs.iowadnr.gov/bionet/Fish/Species/153>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?speciesID=685>

**Eastern Mosquitofish** – Florida Museum of Natural History: <https://www.floridamuseum.ufl.edu/discover-fish/florida-fishes-gallery/eastern-mosquitofish/>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=849>

**Emerald Shiner** – Texas State University, San Marcos, Department of Biology, Freshwater Fishes: <http://txstate.fishesoftexas.org/notropis%20atherinoides.htm>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=582>; Vermont Fish and Wildlife Department: <https://vtfishandwildlife.com/sites/fishandwildlife/files/documents/Fish/baitfish/emerald-shiner.pdf>

**Eurasian Ruffe** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=7>

**Eurasian Rudd** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=648>

**Fallfish** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=650>

**Fathead Minnow** – Missouri Department of Conservation: <https://fieldguide.mt.gov/speciesDetail.aspx?elcode=AFCJB32020>; Montana Natural Heritage Program and Montana Fish, Wildlife and Parks, Montana Field Guide: <https://FieldGuide.mt.gov/speciesDetail.aspx?elcode=AFCJB32020>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=621>

**Fourspine Stickleback** – Connecticut Department of Energy and Environmental Protection: <https://portal.ct.gov/DEEP/Fishing/Freshwater/Freshwater–Fishes–of–Connecticut/Fourspine–Stickleback>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=700>

**Gizzard Shad** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=492>

**Grass Carp** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=514>

**Greenside Darter** – The Regents of the University of Michigan: [https://www.biokids.umich.edu/critters/Etheostoma\\_blennioides/](https://www.biokids.umich.edu/critters/Etheostoma_blennioides/); United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=808>

## Sources

**Golden Shiner** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=579>; Vermont Fish and Wildlife Department: <https://vtfishandwildlife.com/sites/fishandwildlife/files/documents/Fish/baitfish/golden-shiner.pdf>

**Iowa Darter** – Montana Natural Heritage Program and Montana Fish, Wildlife and Parks, Montana Field Guide: <https://fieldguide.mt.gov/speciesDetail.aspx?elcode=AFCQC02240>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=812>

**Mimic Shiner** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=614>; Vermont Fish and Wildlife Department: <https://vtfishandwildlife.com/sites/fishandwildlife/files/documents/Fish/baitfish/mimic-shiner.pdf>

**Mummichog** – Chesapeake Bay Program, Field Guide, Fishes: <https://www.chesapeakebay.net/discover/field-guide/entry/mummichog>; Connecticut Department of Energy and Environmental Protection: <https://portal.ct.gov/DEEP/Fishing/Freshwater/Freshwater–Fishes–of–Connecticut/Mummichog>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?speciesID=688>

**Northern Snakehead** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=2265>

**Redside Dace** – Michigan State University, Natural Features Inventory: <https://mnfi.anr.msu.edu/species/description/11307/Clinostomus–elongatus>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=510>

**Rosyside Dace** – South Carolina Department of Natural Resources:

<https://www.dnr.sc.gov/swap/supplemental/freshwaterfish/rosysidedace2015.pdf>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=511>

**Round Goby** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=713>

**Sculpin** – The Fishes of North Carolina: <https://ncfishes.com/sculpin-family-cottidae-diversity-in-north-carolina/>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=502>

**Silver Carp** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=549>

**Tubenose Goby** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=714>

**Western Mosquitofish** – The University of California, Agricultural and Natural Resources: <https://calfish.ucdavis.edu/species/?uid=109&ds=241>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=846>

**White Perch** – United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=777>

**White Sucker** – United States Department of Agriculture, Forest Service: <https://www.fs.usda.gov/detail/npnht/learningcenter/?cid=fseprd911005>; United States Geological Survey, Nonindigenous Aquatic Species Database: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=346>

## Image Credits

Cover: iStock/Vidya Baduru. Photo of Fish Scales. <https://www.istockphoto.com/photo/closeup-of-the-patterns-as-seen-on-the-body-of-a-fish-gm2149526488-570748721?searchscope=image%2Cfilm>

Page 2: iStock/Serhii Sobolevskyi. Photo of fishing rod and reel. <https://www.istockphoto.com/photo/fisherman-with-rod-spinning-reel-on-the-river-bank-fishing-for-pike-perch-carp-fog-gm1160012188-317395184>

Page 3: iStock/Aleksandr Yurkevich. Photo of perch with bait. <https://www.istockphoto.com/photo/perch-fish-with-bait-vortu-on-a-hook-on-a-fishing-line-sunshine-glint-bright-coloring-gm1381017254-442948010?searchscope=image%2Cfilm>

Page 5: Ashley Wilmont. Photo of New Zealand Mudsnail on a boot  
Pages 6 & 7: iStock/Appfind. Photo of small freshwater fish

<https://www.istockphoto.com/photo/small-fresh-water-fish-gm1308551378-398498047>

Page 10: iStock/slowmotiongli. Photo of Eastern Ruffe. <https://www.istockphoto.com/photo/ruffe-gymnocephalus-cernua-gm1251524889-365275926?clarity=false>

Alewife: Shawn Good/Vermont Fish & Wildlife Department.  
An alewife from Lake Champlain. Copyrighted, All Rights Reserved - Used by Permission. <https://www.fws.gov/media/alewife-lake-champlain>

Banded Darter (female): Uland Thomas – NANFA. Photo of breeding male Banded Darter. United States Geological Survey, Non-indigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=818>

Banded Darter (Male): Uland Thomas – NANFA. Photo of breeding male Banded Darter. United States Geological Survey, Non-indigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=818>

Bighead Carp, juvenile: United States Fish and Wildlife Service. Photo of Bighead Carp. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=551>

Black Carp, juvenile: United States Geological Survey. Columbia Environmental Research Center. Photo of Black Carp.  
<https://www.usgs.gov/media/images/black-and-grass-carp>

Black Bullhead: Chernilevsky, G. (n.d.). Photo of Black Bullhead Catfish. Wikimedia Commons. Licensing:  
<https://creativecommons.org/licenses/by-sa/4.0/>

Bluegill, juvenile: Weigt, L. (2011). Photo of juvenile Bluegill. Smithsonian Environmental Research Institute. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by/2.0/deed.en>

Brook Stickleback: Lethaby, M. (n.d.). Photo of Brook Stickleback. Pennsylvania Sea Grant

Common Carp: iStock/ AleksandarDickov. Photo of Common Carp.  
<https://www.istockphoto.com/photo/carp-gm182851812-13650336>

Creek Chub: Burkhead, N. M. (n.d.). Photo of Creek Chub. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/FactSheet.aspx?speciesID=649>

Eastern Banded Killifish (female): Roy, Z. (2024). Photo of Eastern Banded Killifish. iNaturalist observation: <https://www.inaturalist.org/observations/197490990>. Accessed on October 23, 2024. Licensing: <https://creativecommons.org/licenses/by-nc/4.0/>

Eastern Banded Killifish (male): Aguilar, R. (n.d.). Photo of Eastern Banded Killifish (RR\_08-12-19\_0560). Smithsonian Environmental Research Center. Flickr. Licensing:  
<https://creativecommons.org/licenses/by/2.0/>

Eastern Mosquitofish (female): MarshBunny. (n.d.). Photo of female Eastern Mosquitofish in a Central Floridian Pond. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by-sa/4.0/>

Eastern Mosquitofish (male): Longbottom, W. (2024). Photo of male Eastern Mosquitofish. iNaturalist observation: <https://www.inaturalist.org/observations/248530168>. Accessed on October 23, 2024. Licensing:  
<https://creativecommons.org/licenses/by-nc/4.0/>

## Image Credits

Emerald Shiner: United States Fish and Wildlife Service. (n.d.).  
Photo of Emerald Shiner. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=582>

Eurasian Ruffe: Hunt, T. (n.d.). Photo of Eurasian Ruffe in the Pärnu River, Estonia. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by-sa/3.0/>

European Rudd: Burkhead, N. M. (n.d.). Photo of European Rudd. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=648>

Fallfish: Burkhead, N. M. (n.d.). Photo of Fallfish. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=650>

Fathead Minnow: Aguilar, R. (n.d.). Photo of Fathead Minnow. (44ml SL). Smithsonian Environmental Research Center. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by/2.0/>

Fathead Minnow (breeding male): Simon, J. (2024). Photo of breeding male Fathead Minnow. iNaturalist observation: <https://www.inaturalist.org/observations/229545369>  
Accessed on October 23, 2024. Licensing: <https://creativecommons.org/licenses/by-nc/4.0/>

Fourspine Stickleback: Gatreau, M. (n.d.). Photo of Fourspine Stickleback. University of New Brunswick, invasive.org. Licensing: <https://creativecommons.org/licenses/by-nc/3.0/us/>

Gizzard Shad: Randall, Z. (n.d.). Photo of Gizzard Shad. Florida Museum of Natural History. Copyright information: <https://www.floridamuseum.ufl.edu/about/copyright/>

Golden Shiner: FredlyFish4. (n.d.). Photo of Golden Shiner in Tyson Research Center, Missouri. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by-sa/3.0/>

Grass Carp: United States Fish and Wildlife Service. Photo of Juvenile Grass Carp. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=2948>

Greenside Darter (female): Aguilar, R. (n.d.). Photo of Greenside Darter. Smithsonian Environmental Research Center. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by/2.0/>

Greenside Darter (male): NANFA. Photo of male Greenside Darter. The Fishes of North Carolina. <https://ncfishes.com/freshwater-fishes-of-north-carolina/etheostoma-blennioides/>

Iowa Darter (female): Layne, C. (2024). Photo of female Iowa Darter. iNaturalist observation: <https://www.inaturalist.org/observations/206003459>. Accessed on October 23, 2024 Licensing: <https://creativecommons.org/licenses/by-nc/4.0/>.

Iowa Darter (male): Lethaby, M. (n.d.). Photo of male Iowa Darter. Pennsylvania Sea Grant

Mimic Shiner: United States Fish and Wildlife Service. Photo of Mimic Shiner. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=614>

Mummichog (female): Aguilar, R. (n.d.). Photo of Mummichog. Smithsonian Environmental Research Center. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by/2.0/>

Mummichog (male): Longbottom, W. (2024). Photo of female Mummichog. iNaturalist observation: <https://www.inaturalist.org/observations/230004286>. Accessed on October 23, 2024. Licensing: <https://creativecommons.org/licenses/by-nc/4.0/>

Northern Snakehead: iStock/ Gregory DiSalvo. Photo of Northern Snakehead. <https://www.istockphoto.com/photo/invasive-northern-snakehead-gm2172751772-592506870?searchscope=image%2Cfilm>

Redside Dace: Zell, H. (n.d.). Photo of Redside Dace. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by-sa/3.0/>

## Image Credits

Rosyside Dace: Hagerty, R. (n.d.), United States Fish and Wildlife Service. Photo of Rosyside Dace. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=511>

Round Goby: Slagle, Z. (n.d.), Ohio Department of Natural Resources. Photo of Round Goby. United States Geological Survey, Nonindigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=713>

Sculpin: Matt Tillett. Flickr. Photo of Sculpin. <https://www.flickr.com/photos/mattyfioner/7557837768/>

Silver Carp: U.S. Fish and Wildlife Service. Photo of Common Carp. United States Geological Survey, Non-indigenous Aquatic Species Database. Public Domain: <https://nas.er.usgs.gov/queries/factsheet.aspx?speciesID=549>

Tubenose Goby: Lethaby, M. (n.d.). Photo of Tubenose Goby. Pennsylvania Sea Grant

Western Mosquitofish (female): Tobler, M. (2024). Photo of female Western Mosquitofish. iNaturalist observation: <https://www.inaturalist.org/observations/228673840>. Accessed on October 23, 2024

Western Mosquitofish (male): Tobler, M. (2024). Photo of male Western Mosquitofish. iNaturalist observation: <https://www.inaturalist.org/observations/228673840>. Accessed on October 23, 2024

White Perch: Aguilar, R. (n.d.). Photo of White Perch (91mm SL). Smithsonian Environmental Research Center. Wikimedia Commons. Licensing: <https://creativecommons.org/licenses/by/2.0/>

White Sucker: The Fishes of North Carolina. Photo of White Sucker. <https://ncfishes.com/wp-content/uploads/2021/04/Catostomus-commersonii-white.png>

# Additional Resources

## Fact Sheets

Pennsylvania Sea Grant has developed numerous fact sheets designed for anyone interested in learning more about aquatic invasive species (AIS) and the threats they pose. We've also developed tip sheets, specific to activities known to increase the spread of AIS, with recommendations to help stop the spread of aquatic invaders.

## Guides, Trainings, and Videos

### **STOP Aquatic Invasive Species With the “PA AIS” App**

Aquatic invasive species (AIS) and the “PA AIS” app are featured in this webinar, designed to help users download and navigate through the app to identify and report, in real time, sightings of AIS in Pennsylvania.

### **Pennsylvania Field Guide to Aquatic Invasive Species**

An in-depth guide to aquatic invasive species found in Pennsylvania. The guide includes tips to identify, report, and preventing the spread of aquatic invasive species.


### **Aquatic Invasive Species Video Series**

These videos begin with Aquatic Invasive Species (AIS) “101” and move the audience through the steps to identify, report, and to prevent the spread of AIS.

**Scan Now!** To access AIS resources. Or visit us online at  
<https://seagrant.psu.edu/aquatic-invasive-species/>



# Six Major Watershed Basins



Centimeter Specimen Ruler



## Measurement Conversion

1 centimeter = 0.39 inches

2.5 centimeters = 1 inch

25.4 millimeters = 1 inch

30.5 centimeters = 1 foot

1 meter = 3.28 feet

0.914 meters = 1 yard



PennState